## The stress-secretion relationship in duodenal ulcer\*


The best available evidence suggests that anxiety and emotional tension stimulate acid-pepsin secretion.¹ Also, the development of an ulcer crater in predisposed individuals, or the aggravation of ulcer symptoms, is often typically preceded by a stressful event or situation. Anxiety in particular seems to play a determining role in the course and prognosis of the disease, as well as its etiology.

## A psychobiologic treatment

To obtain comprehensive relief, many duodenal ulcer patients need more than specific, acid-inhibiting medication—they also need dependable reduction of associated anxiety and emotional tension. With adjunctive Librax you can conveniently give your patient both clinical benefits from a single medication.

The pituitary gland plays a key role in the neurohormonal response to emotional stress, leading to an increase in gastric secretion.





## Do you know me?

I'm the first benzodiazepine used clinically
I've been around for 20 years
I have a quarter century of documented efficacy
I have an excellent safety profile
I have a predictable patient response
I have minimal effect on mental acuity; during therapy, however, caution is advised against driving or use of alcohol.

If my name slips your mind, turn the page.



## LIBRIUM CHLORDIAZEROXIDE HC1/ROCHE

\* NOT VALID IF SUBSTITUTE

Dispense :: 134 - 19/11.

ROCHE

Please see adjacent page for summary of product information.